

Immunization Schedule with Combination Vaccines

EVERY FALL: FLU VACCINE⁶ for anyone 6 months and older

	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	4-6 YEARS
PEDIARIX[®] PROQUAD[®] QUADRACEL[™] or KINRIX[®]	PEDIARIX[®] DTaP, IPV, HepB	PEDIARIX[®] DTaP, IPV, HepB ¹	PEDIARIX[®] DTaP, IPV, HepB	HepA	DTaP	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV
	+	+	+	MMR ⁴			+
	PCV	PCV	PCV	Varicella ⁴			PROQUAD[®] MMRV
	Rotavirus	Rotavirus	Rotavirus ²	PCV ⁵			
	Hib	Hib	Hib ³	Hib ⁵			
PENTACEL^{®5} PROQUAD[®] QUADRACEL[™] or KINRIX[®]	PENTACEL[®] DTaP, IPV, Hib	PENTACEL[®] DTaP, IPV, Hib	PENTACEL[®] DTaP, IPV, Hib	HepA	PENTACEL[®] DTaP, IPV, Hib	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV
	+	+	+	MMR ⁴			+
	PCV	PCV	PCV	Varicella ⁴			PROQUAD[®] MMRV
	Rotavirus	Rotavirus	Rotavirus ²	PCV ⁵			
	HepB	HepB ¹	HepB				
VAXELIS^{™8} PROQUAD[®] QUADRACEL[™] or KINRIX[®]	VAXELIS[™] DTaP, IPV, Hib, HepB	VAXELIS[™] DTaP, IPV, Hib, HepB ¹	VAXELIS[™] DTaP, IPV, Hib ³ , HepB	HepA	DTaP	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV
	+	+	+	MMR ⁴			+
	PCV	PCV	PCV	Varicella ⁴			PROQUAD[®] MMRV
	Rotavirus	Rotavirus	Rotavirus ²	PCV ⁵			
			Hib ⁵				

Make sure the vaccine you administer contains the antigens on the doctor's order. Keep it simple. Stick with the same product.

This is a suggested schedule for VFC providers ordering combination vaccines. For alternatives and details, consult the latest "Recommended Immunization Schedules for persons aged 0-18 years, United States." For more info, visit EZIZ.ORG

- A dose of Hepatitis B vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine.
- The six month dose is not needed if Rotarix[®] was used exclusively for both dose 1 and 2 of the rotavirus vaccine series.
- This six month Hib dose is not indicated if PedvaxHIB[®] is used exclusively for the 2 and 4 month infant doses.
- CDC recommends MMR + Varicella at 12-15 months. Providers can use their discretion whether to use MMRV, however.
- Can be administered as late as 15 months. For more information, consult the Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021.

- Influenza vaccine is available in thimerosal-free options. See California Health and Safety Code § 124172.
- Licensed by FDA for children 4 through 6 years with previous doses of IN-FANRIX[™] or PEDIARIX[™]. ACIP recommends that, whenever feasible, the same manufacturer's DTaP vaccines be used for each dose in the series; however, vaccination should not be deferred because the type of DTaP previously administered is unavailable or unknown. See www.cdc.gov/mmwr/preview/mmwrhtml/mm5739a4.htm.
- Licensed by FDA for children 6 weeks through 4 years of age (prior to the 5th birthday).